



**Lectotype of *Celsia acanthifolia* Pau [family
SCROPHULARIACEAE]**

<http://plants.jstor.org/specimen/bc43694>

<http://www.jstor.org>

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the contributing partner regarding any further use of this work. Partner contact information may be obtained at <http://plants.jstor.org/page/about/plants/PlantsProject.jsp>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact plants@jstor.org.

Lectotype of *Celsia acanthifolia* Pau [family SCROPHULARIACEAE]

| | |
|---------------------|--|
| Herbarium | Institut Botànic de Barcelona (BC), BC43694 |
| Collection | Type Specimens |
| Resource Type | Specimens |
| Collector | P. Font Quer, #566 |
| Collection date | 1927/7/6 |
| Locality | pr. Badú (Atlante rhiphaeo); Hab. in saxosis, solo schistoso |
| Country | Morocco (Morocco) |
| Collection altitude | 1500 meters |
| Identifications | <i>Verbascum faurei</i> (Murb.) Hub.-Mor. subsp. <i>acanthifolium</i> (Pau) Benedí & J.M. Monts.-Martí [family SCROPHULARIACEAE] (stored under name); Verified by C. Benedí, 1998/4 <i>Celsia acanthifolia</i> Pau [family SCROPHULARIACEAE]; Lectotype of <i>Celsia acanthifolia</i> Pau [family SCROPHULARIACEAE]; Verified by C. Benedí, 1996 |
| Data last modified | 2010-10-04 |

<http://plants.jstor.org/specimen/bc43694>



BC-43694

Herbario Nacional de España
Instituto Botánico de Barcelona
Barcelona, España



copyright reserved



FLORISTIC BIODIVERSITY OF N MOROCCO

C. faurii
subsp. *acanthifolia* (Pau) (Pau)
Benedict & S. M. Benoit
Determinavit Dr. C. BENEDI IV, 1998

lecto TYPUS
C. Bened. 1998



(A)
43694

DR. FONT QUER. — ITER MAROCCANUM, 1927

596. CELSIA ACANTHIFOLIA Pau, sp. nov.

Hab. in saxosis, pr. Badá (Atlante rhiphaeo), 1500 m. alt., solo schistoso; 6 jull.

Descr.: Folia Acanthi mollis L., superiora oblonga, sinuato-dentata, floribus longe pedunculatis, pedicellis glanduliferis, racemosis; bracteae lanceolatae brevissimae, calycis lacinae lanceolatae, margine integro, capsula globosa, obtusa calyce triplo longior.

BC-43694



REVISION PARA FLORA-HIBERICA N. Marnier
Verbascum farsii (Ranch.) H. & M.
subsp. acanthifolia (Pau) Bonif. &
J. M. Mon S 1996
C. Bonif.
Dpto. Botánica, Fac. Farmacia, Univ. Barcelona



43694⁽³⁾
DR. FONT QUER. — ITER MAROCCANUM, 1927
566. *CELSIA ACANTHIFOLIA* Pau, sp. nov.
Hab. in saxosis, pr. Badú (Atlante rhiphaeo), 1500 m. alt., solo schistoso; 6 Julii.
Descr.: Folia Acanthi mollis L., superiora oblonga, sinuato-dentata, floribus longe pedunculatis, pedicellis glanduliferis, racemosis; bractae lanceolatae brevissimae, calycis laciniae lanceolatae, margine integro, capsula globosa, obtusa calyce triplo longior.



BC-43694



43694⁽²⁾

DR. FONT QUER. — ITER MAROCCANUM, 1927

CELSIA ACANTHIFOLIA Pan, sp. nov.

Hab. in saxosis, pr. Badù (Atlante rhiphaeo), 1500 m. alt., solo schistoso; 6 julli.

Descr.: Folia Acanthi mollis L., superiora oblonga, sinuato-dentata, floribus longe pedunculatis, pedicellis glanduliferis, racemosis; bracteae lanceolatae brevissimae, calycis lacinae lanceolatae, margine integro, capsula globosa, obtusa calyce triplo longior.